

Reanalyses of *Voyager* Ultraviolet Spectrometer Limits to the EUV and FUV Diffuse Astronomical Flux

Jerry Edelstein

jerrye@ssl.berkeley.edu

Stuart Bowyer

bowyer@ssl.berkeley.edu

Michael Lampton

mlampton@ssl.berkeley.edu

Space Sciences Laboratory, University of California, Berkeley, CA 94720

Received _____; accepted _____

ABSTRACT

We re-examine *Voyager* Ultraviolet Spectrograph data used to establish upper limits to the λ 500–900 Å and λ 900–1100 Å cosmic diffuse background. The measurement of diffuse flux with the *Voyager* UVS data requires complex corrections for noise sources which are far larger than the astronomical signal. In the analyses carried out to date, the upper limits obtained on the diffuse background show statistical anomalies which indicate that substantial systematic errors are present. We detail these anomalies and identify specific problems with the analysis. We derive statistically robust 2σ upper limits for continuum flux of 570 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1} \text{Å}^{-1}$ and for the 1000 Å diffuse line flux of 11,790 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1}$. The true limits may be substantially higher because of unknown systematic uncertainties. The new statistical limits alone are insufficient to support previous conclusions based on the *Voyager* data including work on the character of interstellar dust and estimates of the diffuse extragalactic far UV background as absorbed by intergalactic dust.

Subject headings: diffuse radiation – ultraviolet: ISM

1. Introduction

Measurements of the far ultraviolet (FUV: $\lambda\lambda$ 900–1500 Å) and extreme ultraviolet (EUV: $\lambda\lambda$ 90–900 Å) diffuse background are of considerable interest. The FUV background longer than 1200 Å has been categorized in general, (see Bowyer 1991 for a review) but minimal information exists on the band from λ 900–1200 Å. Only upper limits to the general EUV cosmic background exist (see Bowyer et al 1996 for a review of the EUV background); the lowest limits for the EUV background from 500–900 Å have been reported by Edelstein

et al (1999). Possible contributors to the λ 900–1200 Å background include radiation from the hot component of the interstellar medium (ISM), starlight scattered from interstellar dust, emission from fluorescing molecular hydrogen, resonantly scattered radiation from inflowing neutral components of the warm ISM, resonantly scattered geophysical lines, and flux from the electromagnetic decay of elementary particles created in the early universe.

Holberg (1986) and Murthy et al (1991, 1999) have reported the lowest limits for the FUV 900–1100 Å background. These results were obtained using data obtained with the Ultraviolet Spectrometer on the *Voyager* Spacecraft (Broadfoot et al, 1977). Because of a variety of instrumental and environmental effects, the recorded count rate from the *Voyager* instrument was substantially greater than any astronomical diffuse flux; it is about 40 times larger than the claimed upper limit. In principle, a sufficiently precise determination of a stable background will allow an arbitrarily small signal embedded in this background to be identified. However, for data with very low signal to noise even small systematic errors can overwhelm any possibility of deriving a valid signal. We have identified several difficulties in the data set and/or the analysis of Holberg (1986) and Murthy et al (1991) which suggest that significant systematic errors (as compared to the upper limits obtained) are affecting the results. These difficulties are also relevant to the results of Murthy et al (1999) because they incorporate the data reduction procedures of Holberg (1986). In this paper we discuss these problems, and identify some of the systematic effects which are present.

2. Observations and Analysis

Holberg (1986) used 1,508,198 sec of observation from four separate (but astronomically equivalent) regions to obtain limits on a diffuse astronomical flux over the band from 500–1100 Å of 100 to 400 photons s⁻¹ cm⁻² ster⁻¹ Å⁻¹. Holberg did not conduct a formal statistical error analyses and chose an ad-hoc (albeit reasonably considered) error level to

establish his limits. Murthy et al (1991) used observations from four different regions of the sky to obtain 2- σ upper limits on the diffuse flux from 900–1100 Å ranging from -38 to 500 photons s⁻¹ cm⁻² ster⁻¹ Å⁻¹ (see Table 1). Three of these four observations had integrations of $\sim 230,000$ sec, the fourth provided $\sim 83,000$ sec of data. The reduction of both data sets was similar for the majority of the analysis. The major difference was that Murthy et al considered the statistical contribution from only one of the many possible contributors to the ultimate error and employed an additional assumption regarding the statistical independence of data sets that is discussed hereafter.

An immediate indication that large systematic errors may be present in the analysis by Murthy et al (1991) is provided by the data shown in Table 2 of that reference. Systematics can easily vitiate statistical analyses, and the fact that the largest derived fluxes are negative indicates that systematic oversubtraction may have taken place. In that table the authors provide their best estimates for the background in four view directions. To assess the likelihood that the four quoted data are free of systematic error, we have computed the probability of each measurement being consistent with zero flux with a normal probability function. The results are shown in Table 1. Under the null hypothesis of zero continuum flux and no systematic errors being present, the background-subtracted fluxes should group around zero mean and each would exhibit a reasonable ranking against a normal distribution, say 0.1 to 0.9. A negative two σ deviation is a rare event with a probability of $P = 0.023$. Finding two such events in four independent measurements is much rarer still. The probability of such an ensemble can be found by combining all permutations of two special events among four bins, and multiplying by P^2 which is the a priori probability of two special events occurring. This estimate, $6P^2 = 0.004$, reveals the improbability of the ensemble being governed by the stated random errors.

A different kind of test yields the same conclusion. The chi-square test is an aggregate

measure of the total discrepancy represented by an ensemble of measurements. As such, it is tolerant of some data being highly discrepant when other data in the ensemble are less discrepant than average. The total chi square statistic for the four reported measurements is 9.6, a value reached by bias-free statistics in fewer than 5% of all ensembles. This test rejects the hypothesis that the data are fairly described by their random errors at the 95% confidence level.

We note that if we simply ignore the Murthy et al quoted errors, and consider only the scatter in the fluxes assuming that they are measurement samples of zero flux and that no systematic errors are present, we can estimate their uncertainty from their scatter. In this case the most probable value of their standard deviation is $211 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1} \text{ \AA}^{-1}$, a value that would bring the chi squared of their fit to 4.00 for four sample points. This estimate then gives a statistical 2σ upper limit margin of $420 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1} \text{ \AA}^{-1}$ for these data. It is valid to draw statistical conclusions from the set of these four data points – if the errors are not systematic and are estimated properly. Indeed Lampton (1994) has shown that robust statistical conclusions may be determined from just two Poisson samples, one for combined signal and background and the other for background alone. However, our major concern with these data is the very great danger of systematic effects dominating random effects. The finding that two independent tests each show that the published data set is improbable indicates that substantial systematic errors are likely to be present.

We have re-analyzed the long observation of Holberg (1986) toward the north Galactic pole. We consider the uncertainties for each stage of the data reduction process. In Table 2 we list the count rate, the respective photon count given the integration time, and the uncertainty for each of the signal and noise components within a single instrumental bin located at $1000 \text{ \AA}$ of spectral width of $\sim 10 \text{ \AA}$. These values are typical of the entire data

set from 650 – 1000 Å although larger airglow line contributions exist about the Ly- γ and Ly- β spectral regions. We note that the *Voyager* data system registered three counts for each valid photon event (Murthy et al, 1991); the numbers for photon events in our Table 2 are one-third the product of the count rate and the total integration time.

The principal noise components in the *Voyager* data are background counts produced by the on-board radioactive power generator, and instrumental scattering of the interplanetary solar resonance lines of hydrogen and helium. An estimate of the radioactively induced “dark-count” background was obtained by observing a shadowed area of the spacecraft. This spectrum was scaled to fit the 650–900 Å region where no detectable line emission was anticipated, given the instrument’s sensitivity and interstellar absorption. The dark-count background correction corresponds to 11,000 photon events during the observation and represents 58% of the recorded signal. While this background spectrum could, in principle, be stable, Holberg (1986) shows two very long integrations of this background spectrum in which approximately half of the more than 100 spectral channels show variations from spectra to spectra of $\sim 2\%$. These variations are about a factor of two larger than the 1σ statistical counting variation per channel. While we consider 2% of the background count level as an appropriate estimate for the uncertainty in the background, we simply adopt the statistical count rate uncertainty as the minimum background uncertainty for our analysis.

The next manipulation of the data is to remove flux from the strong interplanetary solar resonance lines of H Ly_α , H Ly_β , and He Ly_α which are instrumentally scattered throughout the recorded spectra. The de-scattering correction represents 7,000 photon events during the observation or 37% of the recorded signal. A ground calibration de-scattering matrix was used that redistributes photons among the spectral bins. Errors in either the input spectrum or the de-scattering matrix operator will combine to affect the result. Any background counts that have not been removed from the spectrum will be

redistributed according the matrix operator. Errors within the de-scattering matrix and its interaction with possible errors in background removal are difficult to assess but must be present at some level as demonstrated by the application of the de-scattering technique to an observation of a hot stellar continuum which removed only 93% of the light scattered from above the Lyman limit from the 600–900 Å band stellar flux (Holberg and Barber 1985). We conclude that uncertainties in the de-scattering process are likely to introduce errors that will add to the statistical counting uncertainty. These additional errors cannot be determined from the data presented so we adopt the counting error as the minimum uncertainty for this component.

After the de-scattering process, modeled interplanetary solar resonance lines are subtracted from the spectrum to obtain a residual spectrum from which the limits to an astronomical flux are derived. The uncertainty in the subtraction of interplanetary lines varies across the band, but, excluding the region about Ly- β , the uncertainty due to line flux removal is not a dominant factor. The line modeling correction at 1000 Å represents 500 photon events during the observation or 2.6% of the recorded signal. Systematic error must be present in the line modeling subtraction at some level. For example, Holberg (1986) notes that incomplete removal of the H Ly_{α} line renders these results invalid for $\lambda > 1100$ Å. The magnitude of this uncertainty cannot be determined from the data presented, but must be an addition to the statistical uncertainties.

The residual flux is determined by subtracting the background components from the recorded signal and corresponds to 500 photon events during the observation, or 2.6% of the recorded signal. At a minimum, the uncertainty in the residual flux is dependent on the errors in each of the signal and background components summed in quadrature. The minimum combined uncertainty, considering our discussion above, is 194 photon events (see Table 2). Consequently, the 2σ upper limit to the photon events occurring within a

single channel during the observation, which equals the value of the residual net signal plus twice the standard error, is 888 photon events. We can convert this single channel value to physical continuum units, or $570 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1} \text{ \AA}^{-1}$, given the observation time of $1.5 \times 10^6 \text{ sec}$ and a flux conversion factor determined from the calibrated Voyager spectrum of the white dwarf PG1034+001 of $3.21 \times 10^5 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1} \text{ \AA}^{-1}$ per UVS count per second (Holberg private communication 1997).

To determine the 2σ upper limit to a diffuse monochromatic emission line occurring in the spectra we consider that the line profile characteristic width is three channels wide, or $\sim 30 \text{ \AA}$. If we take the favorable view that errors in three adjacent channels are statistically independent (but see below) then the minimum combined uncertainty in a channel is $194 \times \sqrt{n}$ photon events and the 2σ upper limit is 612 photon events per single channel. Using the flux conversion factor and the $30 \text{ \AA}$ characteristic line width we determine a physical value of the minimum 2σ upper limit to single line emission intensity of 11,790 $\text{photons s}^{-1} \text{ cm}^{-2} \text{ ster}^{-1}$.

We note that the Murthy et al (1991 & 1999) derivation of continuum limits using the 900–1100 $\text{\AA}$ band *Voyager* data followed the same general procedures as Holberg (1986) and consequently the points raised above are also valid for their analysis. In addition these authors assumed that each spectral channel’s errors were uncorrelated. However, the de-scattering process explicitly takes counts from each channel and redistributes them to other channels, which directly violates the assumption of statistical independence. Hence this approach is not valid *prima facie* and its use cannot be justified.

3. Conclusions

Using the published *Voyager* data of Holberg (1986) for observations of the north Galactic pole we estimate a 2σ upper limit for the 1000 Å diffuse line flux of 11,790 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1}$. The limit to continuum flux is 570 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1} \text{Å}^{-1}$. These statistically robust 2σ upper limits are in contrast to the published limits of 100 – 200 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1} \text{Å}^{-1}$ as a 1σ limit (Murthy et al 1991) and 6,000 photons $\text{s}^{-1} \text{cm}^{-2} \text{ster}^{-1}$, with no σ level quoted (Holberg 1986). These new statistical results do not allow for probable systematic effects discussed in the text that are likely raise these limits by some additional unknown amount. Continuum limits based on the new statistical limits alone are greater than the values needed to support a variety of conclusions of Murthy et al (1991) including the derived properties of interstellar dust. We also note that Overduin and Wessin (1997) and Overduin et al (1999) used the Murthey etal (1991) dust results to derive the extragalactic far UV background after absorption by intergalactic dust. These results may now be compromised.

REFERENCES

- Blair, W. P., Vancura, O. Long, K. S. 1995, AJ, 110, 312
- Bowyer, S. 1991, ARA&A, 29, 59
- Bowyer, S., Edelstein, J., Lampton, M., Morales, L., Perez Mercader, J. Gimenez, A. 1996, Astrophysics in the extreme ultraviolet. Proceedings of colloquium no. 152 of the International Astronomical Union; held in Berkeley; California; March 27-30; 1995; Dordrecht: Kluwer Academic Publ.; —c1996; edited by Stuart Bowyer and Roger F. Malina, p.611
- Broadfoot, A. L., et al. 1977, Space Science Reviews, 21, 183 611
- Edelstein, J., Bowyer, S., Korpela, E., Lampton, M., Trapero, J., Gomez, J.F., Morales, C., Orozco, V., 2000, Ap&SS, in press
- Holberg, J. B. Barber, H. B. 1985, ApJ, 292, 16
- Holberg, J. B. 1986, ApJ, 311, 969
- Lampton, M. 1994, ApJ, 436, 784
- Murthy, J. , Henry, R. C. Holberg, J. B. 1991, ApJ, 383, 198
- Murthy, J. , Henry, R. C. Holberg, J. B. 1994, ApJ, 428, 233
- Murthy, J. , Hall, D. , Earl, M. , Henry, R. C. Holberg, J. B. 1999, ApJ, 522, 904
- Overduin, J. M. Wesson, P. S. 1997, ApJ, 483, 77
- Overduin, J. M., Seahra, S. S., Duley, W. W. Wesson, P. S. 1999, A&A, 349, 317

Acknowledgements We acknowledge interesting discussions with Richard Henry.

Table 1: Probability of the Reported Fluxes Being Consistent with Zero

	Reported Flux	Probability	2σ Upper Limit
	CU		CU
Target A	-264 $\pm$ 132	0.023	0
Target B	48 $\pm$ 227	0.6	502
Target C	-72 $\pm$ 138	0.28	200
Target D	-320 $\pm$ 141	0.013	-38
Total Ensemble Permutation probability:		0.004	
Total Ensemble Chi-square probability:		<0.05	

Note: CU = photons s⁻¹ cm⁻² ster⁻¹ Å⁻¹

Table 2: *Voyager* Signal Components per channel

Component	Counting Rate	Photon Events	Minimum Uncertainty ($1\ \sigma$)
Recorded Signal	0.038	19,000	138
Background Components			
Dark-Count	0.022	11,000	105
Instrumental Scattering of Geocoronal Lines	0.014	7,000	84
Interplanetary H Ly_β & H Ly_γ	0.001	500	22
Signal minus Background	0.001	500	
Combined Uncertainty			194
$2\ \sigma$ Upper Limit			888